

Brilliant Violet 605™ anti-human CD193 (CCR3)

Catalog # / 2153575 / 25 tests
Size: 2153580 / 100 tests

Clone: 5E8


Isotype: Mouse IgG2b, κ

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 605™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 605™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).


Concentration: Lot-specific



Human peripheral blood leukocytes were stained with CD16 APC and CCR3 (clone 5E8) Brilliant Violet 605™ (top) or mouse IgG2b Brilliant Violet 605™ isotype control (bottom). Data shown was generated from granulocyte population gating.

Applications:

Applications: Flow Cytometry



Recommended Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 605™ excites at 405 nm and emits at 603 nm. The bandpass filter 610/20 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 605™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes:

Additional reported applications (for the relevant formats) include: The 5E8 antibody is useful for immunofluorescent staining and flow cytometric analysis of CCR3 expression.

It has been observed that the 5E8 antibody clone can interact with PE/Cy7 antibody conjugates during multi-color staining, potentially leading to unwanted staining. This interaction can be resolved by sequentially staining with the 5E8 antibody first and then followed by the PE/Cy7 conjugate of interest.

Application References:

1. Beauvillian C, *et al.* 2011. *Blood* 117:1196. [PubMed](#)

Description: CD193, also known as CC-chemokine receptor 3 (CCR3), CC CKR3, MIP1- α receptor like-2, and eotaxin receptor, is a member of the G protein-coupled seven transmembrane receptors family. It binds to the CC chemokines eotaxin, eotaxin-2, and eotaxin-3 with high affinity. CCR3 has also been reported to bind RANTES, MCP-3, and MCP-4 with low affinity. CCR3 receptor is expressed on human eosinophils, basophils, mast cells, mononuclear phagocytes, platelets, CD34⁺ hematopoietic progenitor cells, Th2-like lymphocytes, and keratinocytes. CCR3 is thought to play a role in allergic diseases such as bronchial asthma and allergic rhinitis. CCR3 is a co-receptor for HIV-1 and HIV-2, and the binding of eotaxin with CCR3 has been shown to inhibit HIV infection in some cell types.

Antigen
References:

1. Gerard W, *et al.* 1996. *J. Exp. Med.* 183:2437.
2. Uguccioni C, *et al.* 1997. *J. Clin. Invest.* 100:1137.
3. Sallusto F, *et al.* 1997. *Science.* 277:2005.
4. Loetscher P, *et a*