

**PE/Cy7 anti-human CD154**

**Catalog # / Size:** 2154160 / 100 tests  
2154155 / 25 tests

**Clone:** 24-31


**Isotype:** Mouse IgG1, κ

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Concentration:** Lot-specific



PMA+ionomycin-stimulated (6 hours) human peripheral blood lymphocytes were stained with CD154 (clone 24-31) PE/Cy7 (filled histogram) or mouse IgG1, κ PE/Cy7 isotype control (open histogram).

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** Additional reported applications (for the relevant formats) include: immunofluorescence microscopy<sup>1,3</sup> and blocking of T cell-dependent B cell differentiation<sup>1,2,4,5</sup>. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 310812). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 310828) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/microg).

- Application References:**
1. Brams P, *et al.* 2001. *Int. Immunopharmacol.* 1:277. (Block, IF)
  2. Rushworth SA, *et al.* 2002. *Transplantation* 73:635. (Block)
  3. Berner B, *et al.* 2000. *Ann. Rheum. Dis.* 59:190. (IF)
  4. Nordström T, *et al.* 2006. *J. Leukocyte Biol.* 79:319. (Block)
  5. Zhang AL, *et al.* 2007. *Blood* doi:10.1182/blood-2007-02-076364. (Block) [PubMed](#)
  6. Kuchen S, *et al.* 2007. *J. Immunol.* 179:5886.
  7. Matus-Nicodermos R, *et al.* 2011. *J. Immunol.* 186:2164. [PubMed](#)

**Description:** CD154 (CD40 ligand) is also known as CD40L, gp39, TRAP and T-BAM. CD40 ligand is a 32-39 kD type II transmembrane glycoprotein. It is a member of the TNF superfamily and is expressed on activated T cells. It has been reported to be important for B cell costimulation following binding of its receptor, CD40. Additionally, binding of CD40L to CD40 on B cells promotes the secretion of immunoglobulin and Ig isotype switching. CD40L is also involved in the regulation of cytokine production by T cells.

- Antigen**
- References:**
1. Najafian N, *et al.* 2003. *Expert Opin. Biol. Ther.* 3:227.
  2. Racke M, *et al.* 2002. *Expert Opin. Ther. Targets.* 6:275.
  3. Ford G, *et al.* 1999. *J. Immunol.* 162:4037.