

**Brilliant Violet 785™ anti-human CD154**

**Catalog # / Size:** 2154205 / 25 tests  
2154210 / 100 tests

**Clone:** 24-31


**Isotype:** Mouse IgG1,  $\kappa$

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 785™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 785™ and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

**Concentration:** 0.2



PMA + ionomycin-stimulated (6 hours) human peripheral blood lymphocytes were stained with CD154 (clone 24-31) Brilliant Violet 785™ (filled histogram) or mouse IgG1,  $\kappa$  Brilliant Violet 785™ isotype control (open histogram).

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 5$  microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 785™ excites at 405 nm and emits at 785 nm. The bandpass filter 780/60 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. **Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel.** Refer to your instrument manual or manufacturer for support. Brilliant Violet 785™ is a trademark of Sirigen Group Ltd.

**Application Notes:** Additional reported applications (for the relevant formats) include: immunofluorescence microscopy<sup>1,3</sup> and blocking of T cell-dependent B cell differentiation<sup>1,2,4,5</sup>. The LEAF™ purified antibody (Endotoxin  $<0.1$  EU/ $\mu$ g, Azide-Free, 0.2  $\mu$ m filtered) is recommended for functional assays (Cat. No. 310812). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 310828) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin  $<0.01$  EU/microg).

**Application References:**

1. Brams P, *et al.* 2001. *Int. Immunopharmacol.* 1:277. (Block, IF)
2. Rushworth SA, *et al.* 2002. *Transplantation* 73:635. (Block)
3. Berner B, *et al.* 2000. *Ann. Rheum. Dis.* 59:190. (IF)
4. Nordström T, *et al.* 2006. *J. Leukocyte Biol.* 79:319. (Block)
5. Zhang AL, *et al.* 2007. *Blood* doi:10.1182/blood-2007-02-076364. (Block) [PubMed](#)
6. Kuchen S, *et al.* 2007. *J. Immunol.* 179:5886.
7. Matus-Nicodermos R, *et al.* 2011. *J. Immunol.* 186:2164. [PubMed](#)

**Description:** CD154 (CD40 ligand) is also known as CD40L, gp39, TRAP and T-BAM. CD40 ligand is a 32-39 kD type II transmembrane glycoprotein. It is a member of the TNF superfamily and is expressed on activated T cells. It has been reported to be important for B cell costimulation following binding of its receptor, CD40. Additionally, binding of CD40L to CD40 on B cells promotes the secretion of immunoglobulin and Ig isotype switching. CD40L is also involved in the regulation of cytokine production by T cells.

**Antigen**  
**References:**

1. Najafian N, *et al.* 2003. *Expert Opin. Biol. Ther.* 3:227.
2. Racke M, *et al.* 2002. *Expert Opin. Ther. Targets.* 6:275.
3. Ford G, *et al.* 1999. *J. Immunol.* 162:4037.