

PerCP/Cy5.5 anti-human CD24

Catalog # / Size: 2155575 / 25 tests
2155580 / 100 tests

Clone: ML5

Isotype: Mouse IgG2a, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V CD24.5

Concentration: Lot-specific



Human peripheral lymphocytes were stained with CD19 FITC and CD24 (clone ML5) PerCP/Cy5.5 (top) or mouse IgG2a isotype control PerCP/Cy5.5 (bottom).

Applications:


Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and 564 nm and a maximum emission of 690 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunofluorescence microscopy3.

- Application References:**
- Schlossman S, *et al.* Eds. 1995. Leukocyte Typing V:White Cell Differentiation Antigens. Oxford University Press. New York.
 - McMichael A, *et al.* 1987. Leukocyte Typing III. Oxford University Press. New York.
 - Yang GP, *et al.* 1999. *Nucleic Acids Research* 27:1517. (IF)
 - Kristiansen G, *et al.* 2003. *Clin. Cancer Res.* 9:4906. (FC)



Description: CD24 is a 35-45 kD glycosylphosphatidylinositol (GPI)-linked protein also known as heat stable antigen (HSA), BA-1, Ly-52, and nectadrin. It is expressed on the surface of B cells (but not plasma cells), granulocytes, follicular dendritic cells, and epithelial cells. CD24 may play a role in the regulation of B-cell proliferation

and maturation. CD24 crosslinking induces a Ca^{2+} flux in mature B cells. CD24 has been shown to interact with CD62P (P-selectin).

**Antigen
References:**

1. Schlossman S, *et al.* Eds. 1995. Leukocyte Typing V. Oxford University Press. New York.