

Brilliant Violet 421™ anti-human CD24

Catalog # / Size: 2155610 / 100 tests
2155605 / 25 tests

Clone: ML5

Isotype: Mouse IgG2a, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Workshop Number: V CD24.5

Concentration: Lot-specific




Human peripheral blood lymphocytes stained with CD19 APC and CD24 (clone ML5) Brilliant Violet 421™ (top) or mouse IgG2a, κ Brilliant Violet 421™ isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for the relevant formats) include:
immunofluorescence microscopy³.

Application References:

1. Schlossman S, *et al.* Eds. 1995. Leukocyte Typing V:White Cell Differentiation Antigens. Oxford University Press. New York.
2. McMichael A, *et al.* 1987. Leucocyte Typing III. Oxford University Press. New York.
3. Yang GP, *et al.* 1999. *Nucleic Acids Research* 27:1517. (IF)
4. Kristiansen G, *et al.* 2003. *Clin. Cancer Res.* 9:4906. (FC)

Description: CD24 is a 35-45 kD glycosylphosphatidylinositol (GPI)-linked protein also known as heat stable antigen (HSA), BA-1, Ly-52, and nectadrin. It is expressed on the surface of B cells (but not plasma cells), granulocytes, follicular dendritic cells, and epithelial cells. CD24 may play a role in the regulation of B-cell proliferation and maturation. CD24 crosslinking induces a Ca²⁺ flux in mature B cells. CD24 has been shown to interact with CD62P (P-selectin).

Antigen References:

1. Schlossman S, *et al.* Eds. 1995. Leukocyte Typing V. Oxford University Press. New York.