## **Product Data Sheet**

## APC/Cy7 anti-human CD24

Catalog # / Size: 2155660 / 100 tests

2155655 / 25 tests

Clone:

Isotype: Mouse IgG2a, κ

**Reactivity:** Human

The antibody was purified by affinity **Preparation:** 

> chromatography and conjugated with APC/Cy7 under optimal conditions. The solution is free of unconjugated APC/Cy7

and unconjugated antibody.


Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Workshop Number: V CD24.5

**Concentration:** Lot-specific



Human peripheral blood lymphocytes were stained with CD19 APC and CD24 (clone ML5) APC/Cv7 (top) or mouse IgG2a, k APC/Cy7 isotype control (bottom).

## **Applications:**

**Applications:** Flow Cytometry

Recommended

**Usage:** 

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be


titrated for optimal performance for

each application.

**Application** Notes: Additional reported applications (for the

relevant formats) include:

immunofluorescence microscopy3.



**Application** References:

1. Schlossman S, et al. Eds. 1995. Leukocyte Typing V:White Cell Differentiation

Antigens. Oxford University Press. New York. 2. McMichael A, et al. 1987. Leucocyte Typing III. Oxford University Press. New

York. 3. Yang GP, et al. 1999. Nucleic Acids Research 27:1517. (IF)

4. Kristiansen G, et al. 2003. Clin. Cancer Res. 9:4906. (FC)

**Description:** CD24 is a 35-45 kD glycosylphosphatidylinositol (GPI)-linked protein also known

as heat stable antigen (HSA), BA-1, Ly-52, and nectadrin. It is expressed on the surface of B cells (but not plasma cells), granulocytes, follicular dendritic cells, and epithelial cells. CD24 may play a role in the regulation of B-cell proliferation and maturation. CD24 crosslinking induces a Ca<sup>2+</sup> flux in mature B cells. CD24

has been shown to interact with CD62P (P-selectin).

**Antigen** References: 1. Schlossman S, et al. Eds. 1995. Leukocyte Typing V. Oxford University Press.

New York.

