## **Product Data Sheet**

## PE/Cyanine5 anti-human CD24

**Catalog** # / 2155740 / 100 tests

**Size:** 2155735 / 25 tests

Clone: ML5

**Isotype:** Mouse IgG2a, κ

Immunogen: Native human IgD protein (full

length)

Reactivity: Human, Other

**Preparation:** The antibody was purified by affinity

chromatography and conjugated with

PE/Cyanine5 under optimal

conditions.


**Formulation:** Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA)

Workshop Number: V CD24.5

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with anti-human CD19 FITC and antihuman CD24 (clone ML5)

PE/Cyanine5 (left) or PE/Cyanine5 mouse IgG2a, κ isotype control

(right).

## **Applications:**

**Applications:** Flow Cytometry

Recommended

**Usage:** 

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5  $\mu L$  per million cells in 100  $\mu L$  staining volume or 5  $\mu L$  per 100  $\mu L$  of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes:

Additional reported applications (for the relevant formats) include:

immunofluorescence microscopy<sup>3</sup>.

Application References:

1. Schlossman S, et al. Eds. 1995. Leukocyte Typing V:White Cell Differentiation Antigens. Oxford University Press. New York.

2. McMichael A, et al. 1987. Leucocyte Typing III. Oxford University Press.

New York.

3. Yang GP, et al. 1999. Nucleic Acids Research 27:1517. (IF) 4. Kristiansen G, et al. 2003. Clin. Cancer Res. 9:4906. (FC)

**Description:** CD24 is a 35-45 kD glycosylphosphatidylinositol (GPI)-linked protein also

known as heat stable antigen (HSA), BA-1, Ly-52, and nectadrin. It is expressed on the surface of B cells (but not plasma cells), granulocytes, follicular dendritic cells, and epithelial cells. CD24 may play a role in the regulation of B-cell proliferation and maturation. CD24 crosslinking induces a  $Ca^{2+}$  flux in mature B cells. CD24 has been shown to interact with CD62P

(P-selectin).

Antigen References: 1. Schlossman S, et al. Eds. 1995. Leukocyte Typing V. Oxford University

**References:** Press. New York.