

PE/Dazzle™ 594 anti-human CD137L (4-1BB Ligand)

Catalog # / Size: 2157580 / 100 tests

Clone: 5F4


Isotype: Mouse IgG1, κ

Reactivity: Human, Non-human primate, Other

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human T lymphoblastic leukemia cell line

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

Application Notes: For most successful immunofluorescent staining results, it may be important to maximize signal over background by using a relatively bright fluorochrome-antibody conjugate (Cat. No. 2157520) or by using a high sensitivity, three-layer staining technique (e.g., including a biotinylated anti-mouse IgG second step (Cat. No. 2626515), followed by SAv-PE (Cat. No. 2626020)).

- Application References:**
1. Akiba H, *et al.* 2000. *J. Exp. Med.* 191:375.
 2. Pollak KE, *et al.* 1995. *Eur. J. Immunol.* 25:488.
 3. DeBenedette MA, *et al.* 1997. *J. Immunol.* 158:551.
 4. Goodwin RG, *et al.* 1993. *Eur. J. Immunol.* 23:2631.

Description: 4-1BB ligand, also known as CDw137L, is a 97 kD member of the TNF superfamily mainly expressed on APCs, activated B and T cells. It has been reported to be important in T cell proliferation and cytokine production through interaction with 4-1BB receptor. 4-1BB ligand appears to be able to act as a costimulatory molecule without the engagement of other costimulatory molecules such as CD28.

- Antigen References:**
1. Akiba H, *et al.* 2000. *J. Exp. Med.* 191:375.
 2. Pollak KE, *et al.* 1995. *Eur. J. Immunol.* 25:488.
 3. DeBenedette MA, *et al.* 1997. *J. Immunol.* 158:551.
 4. Goodwin RG, *et al.* 1993. *Eur. J. Immunol.* 23:2631.