Product Data Sheet

Purified anti-human CD10

Catalog # / Size: 2161010 / 100 μg

Clone: HI10a

Isotype: Mouse IgG1, κ

Reactivity: Human

Preparation: The antibody was purified by affinity


chromatography.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide.

Workshop Number: V CD10.7

Concentration: 0.5



Nalm-6 cell line stained with purified HI10a and detected with anti-mouse IgGs-FITC

Applications:

Applications: Other

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of

this reagent is ≤0.5 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each

application.

Application

Additional reported (for the relevant formats) applications include:

Notes: immunohistochemistry⁶.

Application References:

1. Knapp W. 1989. Leucocyte Typing IV. Oxford University Press New York.

2. Barclay N, et al. 1997. The Leucocyte Antigen Facts Book. Academic Press Inc.

San Diego.

3. Schlossman S, et al. Eds. 1995. Leucocyte Typing V. Oxford University Press.

New York.

4. Denny MF, et al. 2010. J. Immunol. 184:3284. PubMed

5. Yoshino N, et al. 2000. Exp. Anim. (Tokyo) 49:97. (FC)

6. Dall'Era MA, et al. 2007. BMC Urol. 7:3. (IHC)

Description: CD10 is a 100 kD neutral endopeptidase and a member of the metalloprotease

family. It is a type II transmembrane protein also known as common acute lymphoblastic leukemia antigen (CALLA), enkephalinase, and neprilysin. CD10 is expressed on B cell precursors, T cell precursors, and neutrophils. CD10 is involved in B cell development and has been shown to bind opioid enkephalins,

bradykinin, angiotensins I and II, and other biologically active peptides.

Antigen

1. Shipp M, et al. 1993. Blood 82:1052.

References: 2. Lu B, et al. 1995. J. Exp. Med. 181:2271.