

PE/Dazzle™ 594 anti-human CD10

Catalog # / Size: 2161140 / 100 tests
2161135 / 25 tests

Clone: HI10a

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V CD10.7

Concentration: Lot-specific



Human peripheral blood granulocytes were stained with CD10 (clone HI10a) PE/Dazzle™ 594 (filled histogram) or mouse IgG1, κ PE/Dazzle™ 594 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

Application Notes: Additional reported (for the relevant formats) applications include: immunohistochemistry⁶.

Application References:

1. Knapp W. 1989. Leucocyte Typing IV. Oxford University Press New York.
2. Barclay N, *et al.* 1997. The Leucocyte Antigen Facts Book. Academic Press Inc. San Diego.
3. Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
4. Denny MF, *et al.* 2010. *J. Immunol.* 184:3284. [PubMed](#)
5. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
6. Dall'Era MA, *et al.* 2007. *BMC Urol.* 7:3. (IHC)

Description: CD10 is a 100 kD neutral endopeptidase and a member of the metalloprotease family. It is a type II transmembrane protein also known as common acute lymphoblastic leukemia antigen (CALLA), enkephalinase, and neprilysin. CD10 is expressed on B cell precursors, T cell precursors, and neutrophils. CD10 is involved in B cell development and has been shown to bind opioid enkephalins, bradykinin, angiotensins I and II, and other biologically active peptides.

Antigen References:

1. Shipp M, *et al.* 1993. *Blood* 82:1052.
2. Lu B, *et al.* 1995. *J. Exp. Med.* 181:2271.

