

APC anti-human CD180 (RP105)

Catalog # / 2164555 / 25 tests
Size: 2164560 / 100 tests

Clone: MHR73-11

Isotype: Mouse IgG1, κ


Immunogen: Human RP105 complexed with mouse MD-1

Reactivity: Human, Non-human primate, Other

Preparation: The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD180 (clone MHR73-11) APC (filled histogram) or mouse IgG1, κ isotype control APC (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μ l per million cells in 100 μ l staining volume or 5 μ l per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: intracellular staining¹, immunoprecipitation¹, immunohistochemistry¹ of acetone-fixed frozen tissue sections, B cell activation¹, protection from apoptosis induced by radiation or dexamethasone, and induction of the costimulatory molecule CD80 (B7-1)¹.

Application References: 1. Miura Y, *et al.* 1998. *Blood* 92:2815.

Description: CD180, also known as RP105, is a 105 kD member of the toll-like receptor (TLR) family and is expressed on most B cells, monocytes/macrophages and dendritic cells. The TLR family's structural characteristics are critically involved in protein-protein signaling interactions leading to the activation of the innate immune system. Additionally, MD-1, a secretory protein, associates with CD180 (RP105) at the cell surface and this complex, in conjunction with TLR4, recognizes bacterial LPS. The human CD180 (RP105) gene maps to 5q12. The MHR73-11 antibody is able to activate B cells and to protect against irradiation- or dexamethasone-induced apoptosis.

Antigen References: 1. Miura Y, *et al.* 1998. *Blood* 92:2815.