

Brilliant Violet 650™ anti-human/mouse/rat CD278 (ICOS)

Catalog # / 2167745 / 25 tests
Size: 2167750 / 100 tests

Clone: C398.4A

Isotype: Hamster IgG


Immunogen: Mouse T cell clone D10.G4.1

Reactivity: Human, Mouse, Non-human primate, Other, Rat

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 650™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 650™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: Lot-specific



PHA-stimulated human peripheral blood lymphocytes (3 days) stained with CD278 (ICOS, clone C398.4A) Brilliant Violet 650™ (filled histogram) or Armenian hamster IgG isotype control Brilliant Violet 650™ (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 650™ excites at 405 nm and emits at 645 nm. The bandpass filter 660/20 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 650™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: The C398.4A antibody is useful for flow cytometric analysis and is able to costimulate T cell activation and proliferation. Additional reported applications (for the relevant formats) include: immunoprecipitation¹ and *in vitro* costimulation of T cell activation^{1,3,4}.

Application
References:

1. Redoglia V, et al. 1996. *Eur. J. Immunol.* 26:2781.
2. Hutloff A, et al. 1999. *Nature* 397:263.
3. Buonfiglio D, et al. 2000. *Eur. J. Immunol.* 30:3463.
4. Coyle AJ, et al. 2

Description:

ICOS, also known as inducible costimulatory molecule and H4, is a 47-57 kD protein. This protein is homologous to the CD28/CTLA-4 proteins. ICOS is expressed on activated T cells and a subset of thymocytes. It is able to costimulate T cells proliferation. In addition, ICOS is involved in humoral immune responses (B cell germinal center formation). The ICOS ligand is B7h/B7RP-1 or B7-H2. ICOS stimulation has been shown to potentiate TCR-mediated IL-4 and IL-10 production and has been proposed to play a role in Th2 cell development.

Antigen
References:

1. Redoglia V, et al. 1996. *Eur. J. Immunol.* 26:2781.
2. Hutloff A, et al. 1999. *Nature* 397:263.
3. Buonfiglio D, et al. 2000. *Eur. J. Immunol.* 30:3463.
4. Coyle AJ, et al. 2000. *Immunity* 13:95.