
Product Data Sheet

PerCP/Cyanine5.5 anti-human IgM

Catalog # / 2172560 / 100 tests
Size: 2172555 / 25 tests

Clone: MHM-88

Isotype: Mouse IgG1, κ

Immunogen: Human Ig cocktail

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PerCP/Cyanine5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cyanine5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific

□ Overnight cultured human peripheral blood lymphocytes stained with MHM-88 PerCP/Cy5.5

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μ l per million cells or 5 μ l per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cyanine5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes: MHM-88 antibody reacts with both soluble and membrane human immunoglobulin M (IgM). It does not react with other Ig isotypes. Additional reported applications (for the relevant formats) include: use as a primary or secondary reagent for ELISA analysis.

Due to the presence of excess soluble IgM in whole blood, which competes for antibody binding, staining for IgM on cells in whole blood is not recommended.

Application References: 1. Perez-Shiyama C, *et al.* 2014. *J Immunol.* 192:5192. [PubMed](#)

Description: IgM is the first immunoglobulin made by B cells in the immune response. Surface IgM is expressed on immature and mature B cells, while IgM heavy (μ) chain is expressed intracellularly in pre-B cells.