Product Data Sheet

PE/Dazzle™ 594 anti-human IgM

Catalog # / Size: 2172650 / 100 tests

2172645 / 25 tests

Clone: MHM-88

Isotype: Mouse IgG1, κ

Immunogen: Human Ig cocktail

Reactivity: Human

Preparation: The antibody was purified by affinity

chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and


unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Overnight cultured human peripheral blood mononuclear cells were stained with CD19 PE and anti-IgM (clone MHM-88) PE/Dazzle™ 594 (top) or mouse IgG1, κ PE/Dazzle™ 594 isotype control (bottom). Data shown is gated on lymphocyte population

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for

each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.


Application Notes:

MHM-88 antibody reacts with both soluble and membrane human immunoglobulin M (IgM). It does not react with other Ig isotypes. Additional reported applications (for the relevant formats) include: use as a primary or secondary reagent for ELISA analysis.

Due to the presence of excess soluble IgM in whole blood, which competes for antibody binding, staining for IgM on

cells in whole blood is not

recommended.



Application References:

1. Perez-Shiyama C, et al. 2014. J Immunol. 192:5192. PubMed

Description:

IgM is the first immunoglobulin made by B cells in the immune response. Surface IgM is expressed on immature and mature B cells, while IgM heavy (μ) chain is expressed intracellularly in pre-B cells.