


Purified anti-human CD177

Catalog # / Size: 2179010 / 100 µg
Clone: MEM-166
Isotype: Mouse IgG1, κ
Immunogen: Human granulocytes
Reactivity: Human
Preparation: The antibody was purified by affinity chromatography.
Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
Concentration: 0.5



Human peripheral blood granulocytes stained with purified MEM-166, followed by anti-mouse IgG FITC

Applications:

Applications: Other

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤ 2.0 microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation, Western blotting⁵, and immunofluorescence⁴.

Application References: 1. Kishimoto T, *et al.* Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. New York.
2. Angelisová P, *et al.* 1999. *Immunobiology* 200:234.

Description: CD177 is also known as neutrophil specific antigen 1, NB1, and polycythemia rubra vera 1. It is a member of the uPAR family and is a GPI-linked cell surface glycoprotein with a molecular weight of 60 kD. CD177 is expressed on granulocytes and bone marrow progenitors (early erythroblasts, megakaryocytes). It is thought to be involved in allogeneic and autoimmune responses to neutrophils.

Antigen References: 1. Leukocyte Typing VII. Mason D, *et al.* (Eds.) Oxford University Press (2002)
2. Kissel K, *et al.* 2001. *Eur. J. Immunol.* 31:1301.
3. Lalezari P, *et al.* 1971. *J. Clin. Invest.* 50:1108.
4. Teme