Product Data Sheet

86

APC anti-human CD52

Catalog # / Size:	2180035 / 25 tests 2180040 / 100 tests	.h.
Clone:	HI186	
Isotype:	Mouse lgG2b, к	Equipped and the cold Number 100 101 102 103
Immunogen:	Human tonsil	
Reactivity:	Human	
Preparation:	The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.	
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	Log Fluoresence Intensity Human peripheral blood lymphocytes stained with HI18
Concentration:	Lot-specific	APC

Applications:

Applications: Flow Cytometry Recommended Each lot of this antibody is quality control tested by immunofluorescent staining Usage: with flow cytometric analysis. Test size products are transitioning from 20 microL to 5 microL per test. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Application Additional reported applications (for the relevant formats) include: Notes: immunohistochemical staining of formalin-fixed paraffin-embedded tissue sections. Application 1. Kishimoto T, et al. Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. **References:** London. **Description:** CD52, also known as Cambridge pathology antigen 1 (CAMPATH-1), is a 25-29 kD glycoprotein containing a large N-linked carbohydrate moiety. The actual molecule of CD52 is only 8-9 kD. It is expressed in the male reproductive tract and on virtually all lymphocytes (T and B cells), as well as macrophages/monocytes, eosinophils, and red cells. CD52 is thought to play a role in carrying and orienting carbohydrates. CD52 is a potent target for complement-mediated lysis and antibody-mediated cellular cytotoxicity and has been used as a depletion target for chronic lymphocytic leukemia (CLL)/lymphoma and immunosuppression. The HI186 antibody is useful for flow cytometry and immunohistochemistry. Antigen 1. Leukocyte Typing VI. Kishimoto T, et al. (Eds.) Garland Publishing Inc. (1997)

- **References:** 2. Xia MQ, *et al.* 1991. *Eur. J. Immunol.* 21:1677.
 - 3. Kirchhoff C, et al. 1993. Mol. Reprod. Dev. 34:8.
 - 4. Xia MQ, *et al.* <

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com