Product Data Sheet

PE/Cy5 anti-human CD52

Catalog # / 2180115 / 25 tests

Size: 2180120 / 100 tests

Clone: HI186

Isotype: Mouse IgG2b, κ **Immunogen:** Human tonsil

Reactivity: Human, Non-human primate, Other

Preparation: The antibody was purified by affinity

chromatography and conjugated with PE/Cy5 under optimal conditions.


Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA)

Workshop Number: **HCDM** listed

Concentration: Lot-specific



Human peripheral blood lymphocytes stained with CD52 (clone HI186) PE/Cy5 (filled histogram) or mouse IgG2b, κ PE/Cy5 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μL per million cells in 100 μL staining volume or 5 μL per 100 μL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application

Notes:

Additional reported applications (for the relevant formats) include: immunohistochemical staining of formalin-fixed paraffin-embedded tissue sections.

Application References:

1. Kishimoto T, et al. Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. London.

Description:

CD52, also known as Cambridge pathology antigen 1 (CAMPATH-1), is a 25-29 kD glycoprotein containing a large N-linked carbohydrate moiety. The actual molecule of CD52 is only 8-9 kD. It is expressed in the male reproductive tract and on virtually all lymphocytes (T and B cells), as well as macrophages/monocytes, eosinophils, and red cells. CD52 is thought to play a role in carrying and orienting carbohydrates. CD52 is a potent target for complement-mediated lysis and antibody-mediated cellular cytotoxicity and has been used as a depletion target for chronic lymphocytic leukemia (CLL)/lymphoma and immunosuppression. The HI186 antibody is useful for flow cytometry and immunohistochemistry.

Antigen References:

- 1. Leukocyte Typing VI. Kishimoto T, *et al.* (Eds.) Garland Publishing Inc. (1997)
- 2. Xia MQ, et al. 1991. Eur. J. Immunol. 21:1677.
- 3. Kirchhoff C, et al. 1993. Mol. Reprod. Dev. 34:8.
- 4. Xia MQ, et al. 1993. Biochem. J. 293:633.