

**Purified anti-human CD3**

**Catalog # / Size:** 2186510 / 100 µg  
2186505 / 25 µg

**Clone:** OKT3


**Isotype:** Mouse IgG2a, κ

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5



Human peripheral blood lymphocytes stained with purified OKT3, followed by anti-mouse IgGs-FITC

**Applications:**

**Applications:** Flow Cytometry, Immunohistochemistry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.25$  microg per million cells in 100 microL volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** The OKT3 monoclonal antibody reacts with an epitope on the epsilon-subunit within the human CD3 complex.

Clone OKT3 can block the binding of clones SK7 and UCHT1.4 The OKT3 antibody is able to induce T cell activation. Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections and activation of T cells. The LEAF™ purified antibody (Endotoxin  $< 0.1$  EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 317304). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 317326) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin  $< 0.01$  EU/microg).

**Application References:**

- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
- Knapp W. 1989. Leucocyte Typing IV. Oxford University Press New York.
- Barclay N, *et al.* 1997. The Leucocyte Antigen Facts Book. Academic Press Inc. San Diego.
- Li B, *et al.* 2005. *Immunology* 116:487.
- Jeong HY, *et al.* 2008. *J. Leucocyte Biol.* 83:755. [PubMed](#)
- Alter G, *et al.* 2008. *J. Virol.* 82:9668. [PubMed](#)
- Manevich-Mendelson E, *et al.* 2009. *Blood* 114:2344. [PubMed](#)
- Pinto JP, *et al.* 2010. *Immunology.* 130:217. [PubMed](#)
- Biggs MJ, *et al.* 2011. *J. R. Soc. Interface.* 8:1462. [PubMed](#)

**Description:** CD3ε is a 20 kD chain of the CD3/T cell receptor (TCR) complex, which is composed of two CD3ε, one CD3γ, one CD3δ, one CD3ζ (CD247), and a T cell

receptor ( $\alpha/\beta$  or  $\gamma/\delta$ ) heterodimer. It is found on all mature T lymphocytes, NK T cells, and some thymocytes. CD3, also known as T3, is a member of the immunoglobulin superfamily that plays a role in antigen recognition, signal transduction, and T cell activation.

**Antigen  
References:**

1. Barclay N, *et al.* 1993. The Leucocyte FactsBook. Academic Press. San Diego.
2. Beverly P, *et al.* 1981. *Eur. J. Immunol.* 11:329.
3. Lanier L, *et al.* 1986. *J. Immunol.* 137:2501.