

PerCP/Cy5.5 anti-human CD270 (HVEM, TR2)

Catalog # / Size: 2194055 / 25 tests
2194060 / 100 tests

Clone: 122

Isotype: Mouse IgG1, κ


Immunogen: Recombinant human HVEM protein

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD270 (clone 122) PerCP/Cy5.5 (filled histogram) or mouse IgG1, κ PerCP/Cy5.5 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

Application Notes: The 122 antibody has been shown to be useful for flow cytometry, Western blot, and ELISA.

Application References: 1. Cheung TC, *et al.* 2010. *J. Immunol.* 185:1949. [PubMed](#)
2. Hobo W, *et al.* 2012. *J Immunol.* 189:39. [PubMed](#).

Description: The 122 antibody recognizes human HVEM also known as herpesvirus entry mediator A, tumor necrosis factor receptor superfamily, member 14, TNFRSF14, and tumor necrosis factor receptor like 2. HVEM, a member of the TNFR superfamily, is a type I transmembrane protein containing 2 TNF receptor domains with a predicted molecular weight of approximately 30 kD. HVEM is widely expressed in blood vessels, brain, heart, kidney, liver, lung, prostate, spleen, thymus and other organs. Resting T cells and naïve and memory B cells express high levels of HVEM as well. In humans, HVEM is not expressed in germinal center B cells. Immature dendritic cells express high levels of HVEM that is downregulated upon maturation. HVEM plays an important role in herpes simplex virus pathogenesis by enhancing entry into cells. Signaling through HVEM activates JNK1, NF- κ B and AP-1 to control gene expression in response to infection or cellular stress and activate the immune response. HVEM binds to LIGHT and has also been shown to associate with several other proteins including TRAF1, TRAF2, TRAF3, TRAF5, B and T lymphocyte associated protein (BTLA), and

estrogen receptor α .

- Antigen**
References:
1. Carfi A, *et al.* 2001. *Molec. Cell* 8:169.
 2. Gonzalez LC, *et al.* 2005. *Proc. Nat. Acad. Sci.* 102:1116.
 3. Kwon BS, *et al.* 1997. *J. Biol. Chem.* 272:13471.
 4. Marsters SA, *et al.*