## **Product Data Sheet**

## **Purified anti-human FOXP3**

Catalog # / Size:	2201005 / 25 μg 2201010 / 100 μg
Clone:	259D
Isotype:	Mouse IgG1, к
Immunogen:	Full-length FOXP3 protein
<b>Reactivity:</b>	Human
Preparation:	The antibody was purified by affinity chromatography.
Formulation:	This antibody is provided in phosphate- buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Concentration:</b>	0.5



Formalin-fixed, paraffin-embedded Cynomolgus kidney was treated with EDTA pH 8.0 using a high pressure cooker prior to staining. Staining was carried out with monoclonal anti-FoxP3 (clone 259D) at 10 microg/ml followed by biotinylated goat anti-mouse an

## **Applications:**

**Applications:** Other

Recommended

Each lot of this antibody is guality control tested by immunofluorescent Usage: staining with intracellular flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is  $\leq 0.5$  microg per 10<sup>6</sup> cells in 100 microL volume. For Western blotting, the suggested working dilution(s) is  $\leq$  5.0 microg/ml in antibodv dilution buffer. It is recommended that the reagent be titrated for optimal performance for each application.

Application Additional reported applications (for the Notes: relevant formats) include: Western blotting1, and immunohistochemical acetone-fixed frozen staining1 of sections and formalin-fixed paraffinembedded sections. The 259D antibody gives strong positivity on paraffin and frozen sections and the antibody stains some epithelial cells. The binding of 206D to FOXP3 can be partially blocked by 259D, but 206D does not show significant blocking effect on 259D bindina.

> NOTE: For flow cytometric staining with this clone, True-Nuclear™ Transcription



Cell extract from HEK293T cells transfected with human FoxP3 cDNA was resolved by electrophoresis, transferred to nitrocellulose, and probed with monoclonal anti-FoxP3 antibody (clone 259D). Proteins were visualized using a goat anti-mouse secondary conju

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

	Factor Buffer Set (Cat. No. <u>424401</u> ) offers improved staining and is highly recommended.
Application References:	<ol> <li>Roncador G, <i>et al.</i> 2005. <i>Eur. J. Immunol.</i> 35:1681.</li> <li>Yang ZZ, <i>et al.</i> 2006. <i>Blood</i> 107:3639.</li> <li>Liu W, <i>et al.</i> 2006. <i>J. Exp. Med.</i> 203:1701.PubMed</li> <li>Bollyky PL, <i>et al.</i> 2007. <i>J. Immunol.</i> 179:744.</li> <li>Bell MP, <i>et al.</i> 2007. <i>J. Immunol.</i> 179:1893.</li> <li>Tran DQ, <i>et al.</i> 2007. <i>Blood</i> doi:10.1182/blood-2007-06-094656. PubMed</li> <li>Gao Q, <i>et al.</i> 2008. <i>Blood</i> 111:463.PubMed</li> <li>Zheng Y, <i>et al.</i> 2008. <i>J. Immunol.</i> 181:1683. PubMed</li> <li>Zheng Y, <i>et al.</i> 2008. <i>J. Immunol.</i> 181:1683. PubMed</li> <li>Kavanagh B, <i>et al.</i> 2008. <i>Blood</i> 112:287. PubMed</li> <li>Kavanagh B, <i>et al.</i> 2009. <i>Clin Cancer Res.</i> 15:1931. PubMed</li> <li>Nevala WK, <i>et al.</i> 2009. <i>Clin Cancer Res.</i> 15:1931. PubMed</li> <li>Nigam P, <i>et al.</i> 2010. <i>J. Immunol.</i> 184:1690. PubMed</li> <li>Kmieciak M, <i>et al.</i> 2009. <i>J. Transl. Med.</i> 7:89. (ICFC) PubMed</li> <li>Kanghaven S, <i>et al.</i> 2009. <i>Ann Rheum Dis.</i> 68:1908. PubMed</li> <li>Raghaven S, <i>et al.</i> 2014. <i>Cancer Immunol Res.</i> 2:632. PubMed</li> </ol>

**Description:** FOXP3 is a 50-55 kD transcription factor, also known as Forkhead box protein P3, Scurfin, JM2, or IPEX. It is proposed to be a master regulatory gene and more specific marker of T regulatory cells than most cell surface markers (such as CD4 and CD25). Transduced expression of FOXP3 in CD4<sup>+</sup>/CD25<sup>-</sup> cells has been shown to induce GITR, CD103, and CTLA4 and impart a T regulatory cell phenotype. FOXP3 is mutated in X-linked autoimmunity-allergic dysregulation syndrome (XLAAD or IPEX) in humans and in "scurfy" mice. Overexpression of FOXP3 has been shown to lead to a hypoactive immune state suggesting that this transcriptional factor is a central regulator of T cell activity. In human, unlike in mouse, two isoforms of FOXP3 have been reported: one (FOXP3) corresponding to the canonical full-length sequence; the other (FOXP3 δ2) lacking exon 2. The 259D antibody recognizes human FOXP3 epitope in the region of amino acids 105-235.

Antigen 1. Hori S, *et al.* 2003. *Science* 299:1057. References: