SONY

Product Data Sheet

Pacific Blue™ anti-human CD206 (MMR)

Catalog # / Size: 2205755 / 25 tests
2205760 / 100 tests

Clone: 15-2

Isotype: Mouse IgG1, κ

Immunogen: Purified human mannose receptor

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Pacific Blue™ under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA)

Workshop Number: HCDM listed

Concentration: Lot-specific



Human peripheral blood monocytes were stimulated with GM-CSF for 3 days, then stained with CD206 (MMR) (clone 15-2) Pacific Blue™ (filled histogram), or mouse IgG1, κ Pacific Blue™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µL per million cells in 100 µL staining volume or 5 µL per 100 µL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* Pacific Blue™ has a maximum emission of 455 nm when it is excited at 405 nm. Prior to using Pacific Blue™ conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

Application Notes: The 15-2 antibody blocks the interaction of MMR with its ligand, and inhibits mannose receptor-mediated degradation of t-PA by macrophages. Additional reported applications of this antibody (for the relevant formats) include: Western blotting[1], blocking of ligand binding[1,2], immunofluorescence[3], and immunohistochemical staining of acetone-fixed frozen tissue sections[1].



GM-CSF stimulated (3 days) human peripheral blood monocytes were stained with CD206 (clone 15-2) Brilliant Violet 510™ (filled histogram) or mouse IgG1

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.
Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

Application References:

1. Noorman F, *et al.* 1997. *J. Leukocyte Biol.* 61:63. (WB, IHC, Block)
2. Barrett-Bergshoeff M, *et al.* 1997. *Thromb Haemost.* 77:718. (Block)
3. Kato M*, et al*. 2007. *J. Immunol*. 179:6052. (IF)

Description: Macrophage mannose receptor (MMR) is a 162-175 kD type I membrane protein also known as CD206, MRC1, or mannose receptor (MR). It is a pattern recognition receptor (PRR) that belongs to C-type lectin superfamily. MMR is expressed on macrophages, dendritic cells, and hepatic or lymphatic endothelial cells, but not on monocytes. MMR recognizes a range of microbial carbohydrates bearing mannose, fucose, or N-acetyl glucosamine. MMR mediates endocytosis and phagocytosis, induces activation of macrophages and antigen presentation, plays an important role in host defense, and provides a link between innate and adaptive immunity.

Antigen References:

1. Mason D, *et al.* Eds. 2002. Leukocyte Typing VII. Oxford University Press. p303
2. Wileman TE, *et al.* 1986. *P. Natl. Acad. Sci. USA* 83:2501.
3. Apostolopoulos V and McKenzie IF. 2001. *Curr. Mol. Med.* 1:469.
4. Le Cabec V, *et al.* 2005. *J. Leukocyte Biol.* 77:934.
5. Barrett-Bergshoeff M, *et al.* 1997. *Thromb. Haemostatis* 77:718.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.
Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com