

APC anti-human CD140a (PDGFR α)

Catalog # / 2217560 / 100 tests
Size: 2217555 / 25 tests

Clone: 16A1

Isotype: Mouse IgG1, κ


Immunogen: NIH 3T3 cells transfected with human PDGFR α

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human PDGFR α transfected NIH/3T3 cells were stained with CD140a (clone 16A1) APC (filled histogram) or mouse IgG1

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μ l per million cells in 100 μ l staining volume or 5 μ l per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

- Application References:**
1. Gronwald RG, *et al.* 1988. *Proc. Natl. Acad. Sci. USA* 85:3435.
 2. Gilbertson DG, *et al.* 2001. *J. Biol. Chem.* 276:27406.
 3. Seifert RA, *et al.* 1989. *J. Biol. Chem.* 264:8771.
 4. Rupp E, *et al.* 1994. *Eur. J. Biochem.* 225:29.

Description: The 16A1 monoclonal antibody recognizes human CD140a also known as the platelet-derived growth factor receptor, alpha polypeptide, PDGFR2, and PDGFR α . CD140a is a cell surface tyrosine kinase receptor for members of the platelet-derived growth factor family. The identity of the growth factor bound to the receptor determines whether the functional receptor is a homodimer or heterodimer composed of both PDGFR- α and - β . CD140a contains three immunoglobulin-like domains and a tyrosine kinase domain with a predicted molecular weight of approximately 123 kD. CD140a is widely expressed on a variety of mesenchymal-derived cells and has been implicated in the development of some tumors including basal cell carcinoma and gastric stromal cell tumors. Binding of A-chain containing PDGF molecules as well as protease-activated PDGF-C molecules can stimulate cell proliferation. CD140a has been shown to interact with a number of proteins including CRK, Grb2, Grb14, SHP2, and others as integrin β 3, caveolin-1, and nexin sorting molecules. The PDGFR α is heavily phosphorylated on numerous tyrosine residues through both autophosphorylation and ligand-dependent processes. The 16A1 antibody has been shown to be useful for flow cytometric detection of CD140a.

- Antigen**
- References:**
1. Gronwald RG, et al. 1988. *Proc. Natl. Acad. Sci. USA* 85:3435.
 2. Gilbertson DG, et al. 2001. *J. Biol. Chem.* 276:27406.
 3. Seifert RA, et al. 1989. *J. Biol. Chem.* 264:8771.
 4. Rupp E, et al. 1994. *Eur. J. Biochem.* 225:29.