

Brilliant Violet 421™ anti-human CD203c (E-NPP3)

Catalog # / Size: 2223055 / 25 tests
2223060 / 100 tests

Clone: NP4D6

Isotype: Mouse IgG1, κ

Immunogen: HEK-293 cells transfected with human E-NPP3


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Workshop Number: HLDA8

Concentration: Lot-specific




Human peripheral blood lymphocytes were stained with CD193 APC and CD203c (clone NP4D6) Brilliant Violet 421™ (top) or mouse IgG1, κ Brilliant Violet 421™ isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by

U.S. Patent(s), pending patent applications and foreign equivalents.

- Application** 1. Bühring HJ, *et al.* 1999. *Blood* 94:2343.
References: 2. Bühring HJ, *et al.* 2001. *Blood* 97:3303.
3. Platz IJ, *et al.* 2001. *Int. Arch. Allergy Immunol.* 126:335.
4. Charles N, *et al.* 2010. *Nat. Med.* 16:701. (FC) [PubMed](#)
5. Gernez Y, *et al.* 2011. *Int. Arch. Allergy Immunol.* 154:318. (FC) [PubMed](#)
-

Description: CD203c, a transmembrane protein and a member of the ectoenzyme family, is involved in the hydrolysis of extracellular oligonucleotides, nucleoside phosphates, and NAD (possesses ATPase and ATP pyrophosphatase activity). The molecular weight of CD203c is between 130 and 150 kD under reducing conditions and 270 kD under non-reducing conditions. CD203c is expressed on basophils and mast cells, and is highly expressed on activated basophils. Secretory glands in endometrium and glioma cells are also positive. CD203c is a multifunctional ectoenzyme involved in the clearance of extracellular nucleotides whose substrates include nucleoside triphosphates, nucleoside diphosphates, cAMP, and NAD.

- Antigen** 1. Yano Y, *et al.* 2003. *Int. J. Mol. Med.* 12:763.
References: 2. Andoh K, *et al.* 1999. *Biochim. Biophys. Acta.* 1446:213.