

**PE anti-human CD163**

**Catalog # / Size:** 2232525 / 25 tests  
2232530 / 100 tests

**Clone:** RM3/1

**Isotype:** Mouse IgG1, κ


**Immunogen:** Human monocytes

**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Concentration:** Lot-specific



IL-10 stimulated human peripheral blood monocytes stained with RM3/1 PE

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** Clone RM3/1 binds to domain 9 of CD163. Additional reported applications (for the relevant formats) include: immunofluorescence<sup>7</sup>.

**Application References:**

- Högger P, *et al.* 1998. *J. Immunol.* 161:1883. (FC)
- Zwadlo G, *et al.* 1987. *Exp. Cell Biol.* 55:295. (FC)
- Buechler C, *et al.* 2000. *J. Leukoc. Biol.* 67:97. (FC)
- Puig-Kroger A, *et al.* 2009. *Cancer Res.* 69:9395. (FC) [PubMed](#)
- Madsen M, *et al.* 2004. *J. Biol. Chem.* 279:51561. (FC)
- Jones K, *et al.* 2013. *Clin Cancer Res.* 19:731. (FC) [PubMed](#)
- Stewart DA, *et al.* 2012. *Mol. Cancer Res.* 10:727. (IF)

**Description:** CD163 is a member of the group B scavenger receptor cysteine-rich superfamily, also known as GHI/61, M130, RM3/1, p155, hemoglobin-haptoglobin complex receptor, or macrophage-associated antigen. It is a 134 kD (non-reduced)/155 kD (reduced) glycoprotein primarily expressed on macrophages, Kuffer cells, monocytes, subset of dendritic cells, and a subset of hematopoietic stem/progenitor cells. CD163 binds to haptoglobin-hemoglobin complex and TWEAK, and plays a role in clearing hemoglobin and regulating cytokine production by macrophages. Membrane CD163 can be cleaved by metalloproteinases (MMP), resulting in a soluble form. Elevated serum level of sCD163 has been implicated in many kinds of inflammation diseases.

**Antigen References:**

- Roth J, *et al.* 1994. *Transplantation.* 57:127.
- Van den Heuvel MM, *et al.* 1999. *J. Leukoc. Biol.* 66:858.
- Sulahian TH, *et al.* 2000. *Cytokines* 12:1312.

4. *Fabriek BO, et al.*