
## **Product Data Sheet**

## Alexa Fluor<sup>®</sup> 700 anti-human HLA-DR

Catalog # / Size:	2235070 / 100 μg 2235065 / 25 μg
Clone:	LN3
Isotype:	Mouse IgG2b, κ
Immunogen:	human PBL
<b>Reactivity:</b>	Human
Preparation:	The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 700 under optimal conditions.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.
<b>Concentration:</b>	0.5



Human peripheral blood lymphocytes stained with LN3 Alexa Fluor® 700

## **Applications:**

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. The suggested use of this reagent is $\leq$ 2.0 microg per million cells in 100 microL volume. It is highly recommended that the reagent be titrated for optimal performance for each application.
	* Alexa Fluor® 700 has a maximum emission of 719 nm when it is excited at 633 nm / 635 nm. Prior to using Alexa Fluor® 700 conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.
Application Notes:	Additional reported applications (for the relevant formats) include: immunohistochemical staining1 of frozen sections and formalin-fixed paraffin- embedded sections1, and immunoprecipitation1.
Application References:	<ol> <li>Marder RJ, <i>et al.</i> 1985. <i>Lab. Invest.</i> 52:497.</li> <li>Norton AJ and Isaacson PG. 1987. <i>Am. J. Pathol.</i> 128:225.</li> <li>Hua ZX, <i>et al.</i> 1998. <i>Hum. Pathol.</i> 29(12):1441.</li> </ol>
Description:	The LN3 monoclonal antibody reacts with the HLA-DR antigen, a member of MHC class II molecules. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36 kD $\alpha$ (heavy) chain and a 27 kD $\beta$ (light) chain. It is expressed on B cells, activated T cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4 <sup>+</sup> T cells.
Antigen References:	<ol> <li>Levacher M, <i>et al.</i> 1990. <i>Clin. Exp. Immunol.</i> 81:177.</li> <li>Terstappen L, <i>et al.</i> 1990. <i>J. Leuk. Biol.</i> 48:138.</li> <li>Edwards J, <i>et al.</i> 1985. <i>J. Immunol.</i> 137:490.</li> <li>van Es A, <i>et al.</i></li> </ol>

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com