

PE/Cy5 anti-human TCR V γ 9

Catalog # / 2256620 / 100 tests
Size: 2256615 / 25 tests

Clone: B3

Isotype: Mouse IgG1, κ

Immunogen: Human TLR8-transfected cells


Reactivity: Human, Non-human primate

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Cy5 under optimal conditions. The solution is free of unconjugated PE/Cy5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V CD40.5

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD3 (clone UCHT1) APC and anti-human TCR V γ 9 (clone B3) PE/Cy5 (left) or mouse IgG1, κ PE/Cy5 isotype control (right).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μ l per million cells in 100 μ l staining volume or 5 μ l per 100 μ l of whole blood.

Application Notes: Clone KPL-1 is reported to recognize the tyrosine sulfation consensus motif of PSGL-1¹. Additional reported applications (for the relevant formats) include: Western Blot¹, immunoprecipitation², immunohistochemical staining of acetone-fixed frozen tissue sections and formalin-fixed paraffin embedded tissue sections¹, blocks the recognition of PSGL-1 with P- and L-selectin¹.

Application References:

1. Van Rhijn I, et al. 2003. *Intl. Immunol.* 15:373.
2. Yoshino N, et al. 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)

Description: The V γ 9 TCR is a variant of the TCR γ chain expressed on a subset of γ/δ T cells. V γ 9V δ 2 T lymphocytes, a major γ/δ T cell subset in humans, recognize phosphoantigens, certain tumor cells, and cells treated with aminobisphosphonates. This cell population displays cytolytic activity against various tumor cells. The γ/δ TCR is a heterodimeric TCR complex composed of covalently bound γ and δ chains involved in antigen recognition and the non-covalently associated monomeric proteins CD3 δ , γ , ϵ , and ζ chains.

Antigen References:

1. Scotet E, et al. 2005. *Immunity* 22:71
2. Rincon-Orozco B, et al. 2005. *J. Immunol.* 175:2144