

APC/Cyanine7 anti-human TCR V γ 9

Catalog # / Size: 2256635 / 25 tests
2256640 / 100 tests

Clone: B3


Isotype: Mouse IgG1, κ

Reactivity: Human, Non-human primate

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Cyanine7 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA)

Concentration: Lot-specific



Human peripheral blood mononuclear cells were stained with CD3 FITC and TCR V γ 9 (clone B3) APC/Cyanine7 (left) or mouse IgG1, κ APC/Cyanine7 isotype control (right).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 μ L per million cells in 100 μ L staining volume or 5 μ L per 100 μ L of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

- Application References:**
1. Van Rhijn I, *et al.* 2003. *Intl. Immunol.* 15:373.
 2. Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)

Description: The V γ 9 TCR is a variant of the TCR γ chain expressed on a subset of γ/δ T cells. V γ 9V δ 2 T lymphocytes, a major γ/δ T cell subset in humans, recognize phosphoantigens, certain tumor cells, and cells treated with aminobisphosphonates. This cell population displays cytolytic activity against various tumor cells. The γ/δ TCR is a heterodimeric TCR complex composed of covalently bound γ and δ chains involved in antigen recognition and the non-covalently associated monomeric proteins CD3 δ , γ , ϵ , and ζ chains.

- Antigen References:**
1. Scotet E, *et al.* 2005. *Immunity* 22:71
 2. Rincon-Orozco B, *et al.* 2005. *J. Immunol.* 175:2144