

APC/Fire™ 750 anti-human TCR V62

Catalog # / 2257100 / 100 tests
Size: 2257095 / 25 tests

Clone: B6

Isotype: Mouse IgG1, κ

Immunogen: Modified peptide


Reactivity: Human, Other

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: 750 under optimal conditions.

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD3 Alexa Fluor® 488 and TCR V62 (clone B6) APC/Fire™ 750 (top) or mouse IgG1, κ APC/Fire™ 750 isotype control (bottom).

Applications:


Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.

* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

Application Notes: Annexin V Staining

1. Wash cells twice with cold Staining Buffer and then resuspend cells in Annexin V Binding Buffer at a concentration of 1×10^6 cells/mL.
2. Transfer 100 µL of cell suspension in 5 mL test tube.
3. Add 5 µL of APC/Fire™ 750 Annexin V.
4. Add 10 µL of PI solution (Cat. No. 421301) or 7-AAD (Cat. No. 420403/420404).
5. Gently vortex the cells, and incubate for 15 min at room temperature (25°C), in the dark.
6. Add 400 µL of Annexin V Binding Buffer to each tube. Analyze by flow cytometry.



Whole cell extracts (15 µg protein) from the indicated cell lines were resolved by 4-12% Bis-Tris gel electrophoresis, transferred to a PVDF membrane, and probed with 1.0 µg/mL (1:500 dilution) of purified anti-HMOX1 antibody (clone W19398C) overnight at 4°C. Proteins were visualized by chemiluminescence detection using HRP goat anti-rat IgG antibody (Cat. No. 405405) at 2.0 µg/mL dilution. Direct-Blot™ HRP anti-GAPDH antibody (Cat. No. 607904) was used as a loading control at a 1:50000 dilution (lower). Lane M: Molecular weight marker.

Application
References:

1. Rojas RE, *et al.* 2005. *J. Infect. Dis.* 192:1806.
 2. Correia DV, *et al.* 2011. *Blood* 118:992. (FC) [PubMed](#)
-

Description: The Vδ2 TCR is a variant of the TCR δ chain expressed on a subset of γ/δ T cells. Vγ9Vδ2 T lymphocytes, a major γ/δ T cell subset in humans, recognize phosphoantigens, certain tumor cells, and cells treated with aminobisphosphonates. This cell population displays cytolytic activity against various tumor cells. The γ/δ TCR is an heterodimeric TCR complex composed of covalently bound γ and δ chains involved in antigen recognition and the non-covalently associated monomorphic proteins CD3δ, γ, ε, and ζ chains.

Antigen
References:

1. Scotet E, *et al.* 2005. *Immunity* 22:71.
2. Rincon-Orozco B, *et al.* 2005. *J. Immunol.* 175:2144.