## **Product Data Sheet**

## PE anti-human CD1c

**Catalog #** / 2257525 / 25 tests

**Size:** 2257530 / 100 tests

Clone: L161

**Isotype:** Mouse IgG1, κ

Reactivity: Human

**Preparation:** The antibody was purified by affinity

chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE

and unconjugated antibody.


Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Workshop Number: V T-CD01.18

Concentration: Lot-specific



Human peripheral blood

lymphocytes stained with L161 PE

## **Applications:**

**Applications:** Flow Cytometry

**Recommended** Each lot of this a

mended Each lot of this antibody is quality control tested by immunofluorescentUsage: staining with flow cytometric analysis. Test size products are transitioning

from 20 microL to 5 microL per test. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent

be titrated for optimal performance for each application.

**Application** Additional reported applications (for the relevant formats) include:

**Notes:** immunocytochemical staining1.

**Application** 1. M. del Salamone C, et al. 2001. J. Leukoc. Biol. 70:567.

**References:** 2. de Fraissinette A, et al. 1988. Exp. Hematol. 16:764.

2. Li D, et al. 2012. J. Exp Med. 209:109. PubMed

**Description:** CD1c, also known as R7 or M241, is a 43 kD member of the five CD1 antigens

(CD1a-e) in humans. The CD1 molecules are type I glycoprotein with structural

similarities to MHC class I and are non-covalently associated with  $\beta_2\text{-}$ 

microglobulin, belonging to the Ig superfamily. CD1c is expressed on cortical thymocytes, Langerhans cells, dendritic cells, and a subset of B cells. It has been reported that CD1c is also expressed on mature T cells in a tightly regulated manner. CD1c is involved in antigen-presentation of glycolipids. It

may also act in T cells as an immune regulatory molecule.

Antigen 1. Fainboim LM and del C. Salamone. 2002. J. Biol. Reg. Homeos. Ag. 16:125.

References: 2. M. del Salamone C, et al. 2001. J. Leukocyte Biol. 70:567.

3. Zola H, et al. Eds. 2007. Leukocyte and Stromal Cell Molecules:Th