

**Brilliant Violet 421™ anti-human CD1c**

**Catalog # / Size:** 2257630 / 100 tests  
2257625 / 25 tests

**Clone:** L161

**Isotype:** Mouse IgG1, κ


**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

**Workshop Number:** V T-CD01.18

**Concentration:** Lot-specific




Human peripheral blood lymphocytes were stained with CD19 APC and CD1c (clone L161) Brilliant Violet 421™ (top) or mouse IgG1 Brilliant Violet 421™ isotype control (bottom).

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is ≤5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

**Application Notes:** Additional reported applications (for the relevant formats) include:  
immunocytochemical staining<sup>1</sup>.

**Application References:** 1. M. del Salamone C, *et al.* 2001. *J. Leukoc. Biol.* 70:567.  
2. de Fraissinette A, *et al.* 1988. *Exp. Hematol.* 16:764.  
2. Li D, *et al.* 2012. *J. Exp Med.* 209:109. [PubMed](#)

---

**Description:** CD1c, also known as R7 or M241, is a 43 kD member of the five CD1 antigens (CD1a-e) in humans. The CD1 molecules are type I glycoprotein with structural similarities to MHC class I and are non-covalently associated with  $\beta_2$ -microglobulin, belonging to the Ig superfamily. CD1c is expressed on cortical thymocytes, Langerhans cells, dendritic cells, and a subset of B cells. It has been reported that CD1c is also expressed on mature T cells in a tightly regulated manner. CD1c is involved in antigen-presentation of glycolipids. It may also act in T cells as an immune regulatory molecule.

**Antigen References:** 1. Fainboim LM and del C. Salamone. 2002. *J. Biol. Reg. Homeos. Ag.* 16:125.  
2. M. del Salamone C, *et al.* 2001. *J. Leukocyte Biol.* 70:567.  
3. Zola H, *et al.* Eds. 2007. Leukocyte and Stromal Cell Molecules:Th