

APC/Fire™ 750 anti-human CD1c

Catalog # / 2257725 / 25 tests
Size: 2257730 / 100 tests

Clone: L161

Isotype: Mouse IgG1, κ


Reactivity: Human, Non-human primate, Other

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V T-CD01.18

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD19 PE and CD1c (clone L161) APC/Fire™ 750 (left) or Mouse IgG1, κ APC/Fire™ 750 isotype control (right).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.

* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining on frozen tissue^{4, 5} and formalin-fixed paraffin-embedded immunohistochemical staining⁶.

Application References:

1. del C Salamone M, *et al.* 2001. *J Leukoc Biol.* 70:567.
2. de Fraissinette A, *et al.* 1988. *Exp Hematol.* 16:764.
3. Li D, *et al.* 2012. *J Exp Med.* 209:109. [PubMed](#)
4. Xu C, *et al.* 2010. *Am J Hematol.* 85:539 (IHC-F)
5. Gerlini G, *et al.* 2001. *J Invest Dermatol.* 117:576 (IHC-F)
6. Poposki J, *et al.* 2016. *Clin Exp Allergy* 45:384 (IHC-P) [PubMed](#)

Description: CD1c, also known as R7 or M241, is a 43 kD member of the five CD1 antigens (CD1a-e) in humans. The CD1 molecules are type I glycoprotein with structural similarities to MHC class I and are non-covalently associated with β₂-microglobulin, belonging to the Ig superfamily. CD1c is expressed on cortical thymocytes, Langerhans cells, dendritic cells, and a subset of B cells. It has been reported that CD1c is also expressed on mature T cells in a tightly regulated manner. CD1c is involved in antigen-presentation of glycolipids. It may also act in T cells as an immune regulatory molecule.

- Antigen**
- References:**
1. Fainboim LM and del C. Salamone. 2002. *J. Biol. Reg. Homeos. Ag.* 16:125.
 2. M. del Salamone C, *et al.* 2001. *J. Leukocyte Biol.* 70:567.
 3. Zola H, *et al.* Eds. 2007. *Leukocyte and Stromal Cell Molecules: The CD Markers.* P42.