

## **Product Data Sheet**

## FITC anti-human CD319 (CRACC)

Catalog # / Size:	2259090 / 100 tests 2259085 / 25 tests
Clone:	162.1
Isotype:	Mouse lgG2b, κ
Immunogen:	CRACC-Human IgG1 fusion protein
<b>Reactivity:</b>	Cynomolgus,Human
Preparation:	The antibody was purified by affinity chromatography and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
<b>Concentration:</b>	Lot-specific



Human peripheral blood lymphocytes were stained with CD3 PE and CD319 (clone 162.1) FITC (top) or mouse IgG2b



App	licatio	ns:
-----	---------	-----

Applications: Flow Cytometry

Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 $\mu$ l per million cells or 5 $\mu$ l per 100 $\mu$ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.	Motited InCOh in feature

Application	1. Bouchon A, <i>et al.</i> 2001. <i>J. Immunol.</i> 167:551721.
<b>References:</b>	2. Tassi I and Colonna M. 2005. <i>J. Immunol.</i> 175:79968002.
	3. Veillette A. 2006. <i>Immunol.</i> Rev. 214:2234.

Description:	CD319 is a single-pass type I transmembrane glycoprotein, expressed on NK cells, subsets of mature dendritic cells, activated B cells, and cytotoxic lymphocytes, but not in promyelocytic, B or T cell lines. Expression is highest in the spleen, lymph nodes, and peripheral blood leukocytes, and lowest in bone marrow. Additionally, it is expressed in the small intestine, stomach, appendix, lung, and trachea. CD319 is tyrosine phosphorylated in activated NK cells and is associated with 19 and 39 kD proteins. CD319 has homology with the CD2 family of receptors within the Ig superfamily. Some of the CD2 members stimulate cytotoxicity through the CD319 associated protein.
Antigen	1. Bouchon A, <i>et al.</i> 2001. <i>J. Immunol.</i> 167:551721.

**References:** 2. Tassi I and Colonna M. 2005. *J. Immunol.* 175:79968002.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com