
## **Product Data Sheet**

## APC anti-human CD335 (NKp46)

Catalog # / Size:	2259585 / 25 tests 2259590 / 100 tests
Clone:	9E2
Isotype:	Mouse IgG1, κ
Immunogen:	NKp46-Fc fusion protein
<b>Reactivity:</b>	Human
Preparation:	The antibody was purified by affinity chromatography and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).
<b>Concentration:</b>	Lot-specific



Human peripheral blood lymphocytes were stained with CD56 PE and CD335 (clone 9E2) APC (top) or mouse IgG1,  $\kappa$  APC isotype control (bottom).

105 -

## **Applications:**

Applications:	Flow Cytometry	404
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.	Dd v 10 <sup>2</sup> 0 10 <sup>2</sup> 0 10 <sup>2</sup> 0 10 <sup>2</sup> CD56 PE
Application Notes:	Clone 9E2 has been shown to block NK activation through NKp46. <sup>6</sup>	
Application References:	<ol> <li>Nakajima H, <i>et al.</i> 2000. <i>Eur. J. Immunol.</i> 30:3309.</li> <li>Kalberer CP, <i>et al.</i> 2003. <i>Blood</i> 102:127.</li> <li>Chen Y, <i>et al.</i> 2007. <i>J. Immunol.</i> 179:2766.</li> <li>Jarahian M, <i>et al.</i> 2009. <i>J. Virol.</i> 83:8108. <u>PubMed</u></li> <li>Correia DV, <i>et al.</i> 2011. <i>Blood</i> 118:992. (FC) <u>PubMed</u></li> <li>Achdout H. <i>et al.</i> 2010. <i>J. Virol.</i> 84:3993.</li> </ol>	
Description:	CD335, also known as NKp46, is a member of the natural cytotoxicity receptor (NCR) family which triggers cytotoxicity in NK cells. CD335 is directly involved in target cell recognition and lysis, and is exclusively expressed on CD3 <sup>-</sup> CD56 <sup>+</sup> NK cells, suggesting it is a universal marker for NK cells. NKp46, along with NKp30 and NKp44, is referred to as a natural cytoxicity receptor (NCR) and plays a very important role in killing virus-infected tumor cells and MHC-class I-unprotected cells.	
Antigen References:	1. Mandelboim O and Porgador A. 2001. <i>Int. J. Biochem. Cell Biol.</i> 33:1147. 2. Nakajima H, <i>et al.</i> 2000. <i>Eur. J. Immunol.</i> 30:3309.	
research use only. N	lot for diagnostic use. Not for resale. Sony Bi	iotechnology Inc. will not be held respor

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com 3. Sivori S. 1999. Eur. J. Immunol. 29:1656.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com