
## **Product Data Sheet**

## APC anti-human CD79a (Igα)

Catalog # / Size:	2267530 / 100 tests 2267525 / 25 tests	ъг
Clone:	HM47	
Isotype:	Mouse IgG1, κ	3
<b>Reactivity:</b>	Human	CD19 FITC
Preparation:	The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.	
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	Huma
Workshop Number:	V cB017	lympl CD19
<b>Concentration:</b>	Lot-specific	stain



Human peripheral blood ymphocytes surface stained with CD19 (HIB19) FITC and intracellular stained with HM47 APC

## **Applications:**

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. <b>Test size products</b> <b>are transitioning from 20 microL to 5 microL per test</b> . Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Application References:	1. Mason DY, <i>et al.</i> 1991. J. Immunol. 147:2474 2. Bhargava P, <i>et al.</i> 2007. <i>Am. J. Clin. Pathol.</i> 128:306
Description:	CD79a is a 47 kD type I integral membrane protein, also known as mb-1 or Iga. It is a member of the Ig superfamily and disulphide-associated with CD79b (B29). The interaction of CD79a/CD79b heterodimer with B cell suface Ig forms B cell antigen complex. CD79a is expressed in B cells from early pre-B to plasma cell stage. It has been shown that CD79a is also weakly expressed in some precursors of T- and myeloid cells. CD79 mediates the transport of IgM to B cell surface and transduces signals initiated by BCR aggregation.
Antigen References:	<ol> <li>Zola Heddy, <i>et al.</i> Eds. 2007. Leukocyte and Stromal Cell Molecules:The CD markers. WILEY-LISS</li> <li>Astsaturov IA, <i>et al.</i> 1996. <i>Leukemia</i> 10:769</li> <li>Mson DY, <i>et al.</i> 1995 <i>Blood</i> 86:1453</li> <li>Hashimo</li> </ol>

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com