

APC anti-human CD163

Catalog # / 2268045 / 25 tests
Size: 2268050 / 100 tests

Clone: GHI/61

Isotype: Mouse IgG1, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: VI M38

Concentration: Lot-specific



Human peripheral blood monocytes stained with GHI/61 APC

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Clone GHI/61 binds to domain 7 of CD163. Additional reported applications (for the relevant formats) include: immunocytochemical staining, immunoprecipitation, and western blot.¹

Application References:

1. Pulford K, *et al.* 1992. *Immunology* 75:588. (ICC, IP, WB)
2. Law SK, *et al.* 1993. *Eur. J. Immunol.* 23:2320.
3. Madsen M, *et al.* 2004. *J. Biol. Chem.* 279:51561.
4. Kim WK, *et al.* 2006. *Am. J. Pathol.* 168:822. (FC)
5. Buttari B, *et al.* 2011. *Atherosclerosis*. 215:316. [PubMed](#)

Description: CD163 is a member of the group B scavenger receptor cysteine-rich superfamily, also known as GHI/61, M130, RM3/1, p155, hemoglobin-haptoglobin complex receptor, or macrophage-associated antigen. It is a 134 kD (non-reduced)/155 kD (reduced) glycoprotein primarily expressed on macrophages, Kupffer cells, monocytes, a subset of dendritic cells, and a subset of hematopoietic stem/progenitor cells. CD163 binds to haptoglobin-hemoglobin complex and TWEAK, and plays a role in clearing hemoglobin and regulating cytokine production by macrophages. Membrane CD163 can be cleaved by metalloproteinases (MMP), resulting in a soluble form. Elevated serum level of sCD163 has been implicated in many kinds of inflammatory diseases.

- Antigen** 1. Roth J, et al. 1994 *Transplantation*. 57:127
- References:** 2. Van den Heuvel MM, et al. 1999 *J. Leukoc. Biol.* 66:858
3. Sulahian TH, et al. 2000 *Cytokines* 12:1312
4. Fabrick BO, et al. 20