

PE/Dazzle™ 594 anti-human CD85j (ILT2)

Catalog # / Size: 2268580 / 100 tests
2268575 / 25 tests

Clone: GHI/75

Isotype: Mouse IgG2b, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PE/Dazzle™ 594 under optimal conditions. The solution is free of unconjugated PE/Dazzle™ 594 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V B032

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD85j (Clone: GHI/75) PE/Dazzle™ 594 (filled histogram) or mouse IgG2b, κ PE/Dazzle™ 594 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PE/Dazzle™ 594 has a maximum excitation of 566 nm and a maximum emission of 610 nm.

Application Notes: Additional reported application: Block HLA-G induced TGF-β1 production.

Application References:

- Zola H, *et al.* 2007. Leukocyte and Stromal Cell Molecules: The CD Markers Wiley-Liss A John Wiley & Sons Inc, Publication
- Kirwan SE and Burshtyn DN. 2005. *J. Immunol.* 175:5006

Description: CD85 is a group of Ig superfamily transmembrane glycoproteins called Ig-Like Transcripts (ILTs) or Leukocyte Immunoglobulin-like Receptors (LIRs). CD85j is the 110kD member, known as ILT2, LIR1, or LILRB1, and MIR7. ILT2 structurally has four Ig domains and contains ITIMs in its cytoplasmic tail that provide inhibitory signals by recruiting SHP-1. ILT2 is found on the surface of B cells, plasma cells, dendritic cells, monocytes, subsets of NK and T cells. The ligands of ILT2 include a broad range of HLA-A, -B molecules, some HLA-C and HLA-G molecules, and the human cytomegalovirus UL18 protein.

Antigen References:

- Zola H, *et al.* 2007. Leukocyte and Stromal Cell Molecules: The CD Markers Wiley-Liss A John Wiley & Sons Inc, Publication
- Kirwan SE and Burshtyn DN. 2005. *J. Immunol.* 175:5006