

Brilliant Violet 421™ anti-human CD68

Catalog # / Size: 2269135 / 25 tests
2269140 / 100 tests

Clone: Y1/82A

Isotype: Mouse IgG2b, κ


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Workshop Number: VI MR23

Concentration: Lot-specific



Human peripheral blood monocytes were fixed, permeabilized, and intracellularly stained with CD68 (clone Y1/82A) Brilliant Violet 421™ (filled histogram) or Brilliant Violet 421™ mouse IgG2b, κ isotype control (open histogram).

Applications:

Applications: Intracellular Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.

Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported application: immunohistochemical staining of frozen tissue sections. This clone was tested in-house and does not work on formalin fixed paraffin-embedded (FFPE) tissue.

- Application References:**
1. Doussis IA, et al. 1993. *J. Clin. Pathol.* 46:334.
 2. Davey FR, et al. 1988. *J. Clin. Pathol.* 41:753.
 3. Bushway ME, et al. 2014. *Biol Reprod.* 90(5): 110. (IF) [PubMed](#)

Description: CD68 is a 110 kD glycoprotein, also known as macrosialin, belonging to the sialomucin family. It is closely related to the family of acidic, highly glycosylated lysosomal-associated membrane proteins (LAMPs). CD68 is predominately expressed in cytoplasmic granules of monocytes/macrophages, dendritic cells, and granulocytes. It is one of the useful myeloid cell markers. Further studies have shown that CD68 is also expressed by a subset of hematopoietic progenitors, γ/δ T cells, NK cells, LAK cells, subset of B cells, fibroblasts, and endothelial cells. The biological function of CD68 is still unknown.

Antigen
References:

1. Holness CL and Simmons DL. 1993. *Blood* 81:1607.
2. Gottfried E, et al. 2008. *Scand. J. Immunol.* 67:453.
3. Hameed A, et al. 1994. *Hum. Pathol.* 25:872.