

**APC/Fire™ 750 anti-human FcεRIα**

**Catalog # / Size:** 2273215 / 25 tests  
2273220 / 100 tests

**Clone:** AER-37 (CRA-1)


**Isotype:** Mouse IgG2b, κ

**Reactivity:** Human, Non-human primate, Other

**Preparation:** The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Concentration:** Lot-specific



Human peripheral blood lymphocytes were stained with CD203c FITC and anti-human FcεRIα (clone AER-37 (CRA-1)) APC/Fire™ 750 (left) or Mouse IgG2b, κ APC/Fire™ 750 isotype control (right).

**Applications:**

**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.

\* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

**Application Notes:** Clone AER-37 (CRA-1) has been reported to bind the receptor even in the presence of IgE.<sup>4</sup>

- Application References:**
1. Yamaguchi M, *et al.* 1999. *J. Immunol.* 162:5455.
  2. Suzukawa M, *et al.* 2005. *Int. Immunol.* 17:1249.
  3. Charles N, *et al.* 2010. *Nat. Med.* 16:701. (FC) [PubMed](#)
  4. Yamaguchi M, *et al.* 1999. *J. Immunol.* 162:5455.

**Description:** High affinity IgE receptor (FcεRI) plays a key role in IgE-mediated allergic immune response. FcεRI is a tetrameric receptor complex, which is composed of one α-subunit (FcεRIα), one β-subunit, and two γ-subunits. FcεRIα directly binds IgE with high affinity, while the β- and γ-chains are responsible for mediating intracellular signals. FcεRIα is a 50 kD transmembrane protein with Ig superfamily structure. It is primarily found on mast cells and basophils. Further studies have indicated that FcεRIα is also expressed on many inflammatory cells including cutaneous Langerhans cells, dendritic cells, monocytes of patients with allergic disorders, platelets, bronchial epithelial cells, eosinophils produced in hypereosinophilic syndrome, and neutrophils from allergy-induced asthma patients.

- Antigen**
- References:**
1. Riske F, *et al.* 1991. *J. Biol. Chem.* 266:11245
  2. Gounni AS, *et al.* 2001. *FASEB J.* 15:940.
  3. Maurer D, *et al.* 1996. *J. Immunol.* 157:607
  4. Maurer d, *et al.* 1994. *J. Exp. Med.* 179:745
  5. Campbell AM, *et al.* 1998. *Am. J. Respir. Cell Mol. Biol.* 19:92.