

Brilliant Violet 421™ anti-human CD36

Catalog # / Size: 2281145 / 25 tests
2281150 / 100 tests

Clone: 5-271

Isotype: Mouse IgG2a, κ


Immunogen: Human platelets

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 421™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 421™ and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood platelets were stained with CD36 (clone 5-271) Brilliant Violet 421™ (filled histogram) or mouse IgG2a, κ Brilliant Violet 421™ isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.

Brilliant Violet 421™ excites at 405 nm and emits at 421 nm. The standard bandpass filter 450/50 nm is recommended for detection. Brilliant Violet 421™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

Application Notes: Additional reported applications (for the relevant formats) include: immunofluorescence2.

Application References:
 1. Stelner E, et al. 2006. *J. Cell Sci.* 119:459.
 2. Stewart DA, et al. 2012. *Mol. Cancer Res.* 10:727. (IF)

Description: CD36 is an 85 kD integral membrane glycoprotein, also known as GPIIb, or GPIV. It is expressed on various epithelial and endothelial cells as well as erythrocytes, platelets, macrophages/monocytes and some macrophage-derived dendritic cells. CD36 functions as a scavenger receptor, binding thrombospondin, long chain fatty acids, oxidized LDL, collagen type I, IV, and V as well as apoptotic cells. The 5-271 antibody has been reported to be useful for flow cytometry.

- Antigen**
- References:**
1. Hogg N, *et al.* 1984. *Immunology* 53-753.
 2. Greenwalt DE, *et al.* 1992. *Blood* 80:1105.
 3. Armesilla AL, *et al.* 1994. *J. Biol. Chem.* 269:18985.
 4. Endemann G, *et al.* 1993. *J. Biol. Chem.* 268:11811.