

APC/Fire™ 750 anti-human CD61

Catalog # / Size: 2282105 / 25 tests
2282110 / 100 tests

Clone: VI-PL2


Isotype: Mouse IgG1, κ

Reactivity: Human, Non-human primate, Other

Preparation: The antibody was purified by affinity chromatography and conjugated with APC/Fire™ 750 under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood platelets were stained with CD61 (clone VI-PL2) APC/Fire™ 750 (filled histogram) or mouse IgG1, κ APC/Fire™ 750 isotype control (open histogram).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 µl per million cells in 100 µl staining volume or 5 µl per 100 µl of whole blood.

* APC/Fire™ 750 has a maximum excitation of 650 nm and a maximum emission of 787 nm.

Application Notes: Additional reported applications (for the relevant formats) include: Western blotting and immunohistochemical staining of frozen tissue sections.

- Application References:**
1. Davies J, *et al.* 1989. *J. Cell Biol.* 109:1817.
 2. Roberts M, *et al.* 2004. *Mol. Cell. Biol.* 24:1505.
 3. Ciarlet M, *et al.* 2002. *J. Virol.* 76:1109.

Description: CD61, also known as integrin β3 and glycoprotein IIIa (gpIIIa), is a 90 kD type I integral transmembrane glycoprotein. It is a member of the integrin family, associating with platelet gpIIb (CD41) to form CD41/CD61 complex and with integrin αV (CD51) to form αV/β3 (CD51/CD61) integrin. CD41/CD61 is expressed on platelets and megakaryocytes, and plays a role in platelet activation and aggregation through interaction with fibrinogen, fibronectin, vWF, and other RGD-containing adhesion molecules. CD51/CD61 is expressed on platelets, osteoclasts, fibroblasts, macrophages, and some tumor cells involved in tumor metastasis, and in adenovirus infection through binding to RGD motif in extracellular matrix proteins.

Antigen References:

1. Zola H, *et al.* 2007. *Leukocyte and Stromal Cell Molecules: The CD Markers.*