

**Brilliant Violet 650™ anti-human CD11c**


<b>Catalog # / Size:</b>	2286185 / 25 tests 2286190 / 100 tests
<b>Clone:</b>	Bu15
<b>Isotype:</b>	Mouse IgG1, $\kappa$
<b>Reactivity:</b>	Human
<b>Preparation:</b>	The antibody was purified by affinity chromatography and conjugated with Brilliant Violet 650™ under optimal conditions. The solution is free of unconjugated Brilliant Violet 650™ and unconjugated antibody.
<b>Formulation:</b>	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).
<b>Workshop Number:</b>	V S143
<b>Concentration:</b>	Lot-specific

**Applications:**

- Applications:** Flow Cytometry
- Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5  $\mu$ l per million cells or 5  $\mu$ l per 100  $\mu$ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Brilliant Violet 650™ excites at 405 nm and emits at 645 nm. The bandpass filter 660/20 nm is recommended for detection, although filter optimization may be required depending on other fluorophores used. Be sure to verify that your cytometer configuration and software setup are appropriate for detecting this channel. Refer to your instrument manual or manufacturer for support. Brilliant Violet 650™ is a trademark of Sirigen Group Ltd.

This product is subject to proprietary rights of Sirigen Inc. and is made and sold under license from Sirigen Inc. The purchase of this product conveys to the buyer a non-transferable right to use the purchased product for research purposes only. This product may not be resold or incorporated in any manner into another product for resale. Any use



for therapeutics or diagnostics is strictly prohibited. This product is covered by U.S. Patent(s), pending patent applications and foreign equivalents.

**Application Notes:** Clone Bu15 has a different binding epitope than clone 3.9. The binding of Bu15 with CD11c is divalent cation independent. Additional reported applications (for the relevant formats of this clone) include: inhibition of CD11c mediated adhesion and stimulation of chemokine production by monocytes.

**Application References:**

1. Petty H. 1996. *Immunol. Today* 17:209.
2. Springer T. 1994. *Cell* 76:301.
3. Ihanus E, *et al.* 2007. *Blood* 109:802-810.

---

**Description:** CD11c is a 145-150 kD type I transmembrane glycoprotein also known as integrin  $\alpha_x$  and CR4. CD11c non-covalently associates with integrin  $\beta_2$  (CD18) and is expressed on monocytes/macrophages, dendritic cells, granulocytes, NK cells, and subsets of T and B cells. CD11c has been reported to play a role in adhesion and CTL killing through its interactions with fibrinogen, CD54, and iC3b.

**Antigen References:**

1. Petty H. 1996. *Immunol. Today* 17:209.
2. Springer T. 1994. *Cell* 76:301.
3. Ihanus E, *et al.* 2007. *Blood* 109:802-810.