Alexa Fluor® 647 anti-GFP

Catalog # / Size: 2290030 / 100 tests

2290025 / 25 tests

Clone: FM264G Isotype: Rat IgG2a, κ

Immunogen: TLR9-GFP transfected cell line

Reactivity: Human

Preparation: The antibody was purified by affinity

chromatography, and conjugated with Alexa Fluor® 647 under optimal


conditions.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



GFP transfected CHO cells intracellularly stained with FM264G Alexa Fluor® 647 or isotype control rat IqG2a Alexa Fluor® 647

Applications:

Applications: Flow Cytometry

Recommended Usage:

Each lot of this antibody is quality control tested by intracellular

immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for

optimal performance for each application.

* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at

633nm / 635nm.

Application

1. Chen G, et al. J. Virol. 85:1131. PubMed

References:

2. Luo Y, et al. 2012. J Control Release. 162:28. PubMed

3. Zuo X, et al. 2014. PLoS One. 9:84748. PubMed

Description: Green fluorescent protein (GFP) was originally identified as a protein involved in

bioluminescence, which is from the jellyfish *Aequorea Victoria*. It is widely used as a fluorescent indicator for monitoring gene expression in a variety of cellular systems, including living organisms and fixed tissues. Unlike other bioluminescent reporters, GFP fluoresces without the need for exogenous substrates or cofactors, or other intrinsic or extrinsic proteins, making GFP a useful tool for monitoring gene expression and protein localization *in vivo*. Purified GFP is a 27 kD monomer consisting of 238 amino acids and emits green light (emission maximum at 509

nm) when excited with blue or UV light.

Antigen References:

1. Ishikura H, et al. 2004. Anticancer Res. 24:719.

2. Rizzuto R, et al. 1996. Curr. Biol. 6:183.

3. Chalfie M, et al. 1994. Science 263:802.