## **Product Data Sheet**

## PE anti-human CD44

Catalog # / Size:	2294035 / 25 tests 2294040 / 100 tests	Egetive Cell Number 10 <sup>0</sup> 10 <sup>1</sup> 10 <sup>2</sup> 10 <sup>3</sup> 10 <sup>4</sup>
Clone:	BJ18	
Isotype:	Mouse IgG1, к	
Immunogen:	Normal human PBL	
<b>Reactivity:</b>	Human	
Preparation:	The antibody was purified by affinity chromatography, and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.	
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	Log Fluorescence Intensity Human peripheral blood lymphocytes stained with BJ18 PE
Workshop Number:	VI A034	
<b>Concentration:</b>	Lot-specific	

## **Applications:**

Applications:	Flow Cytometry
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. <b>Test size products are transitioning from 20 microL to 5 microL per test</b> . Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.
Application References:	<ol> <li>Kishimoto T, <i>et al.</i> eds. 1997 <i>Leucocyte Typing VI:White Cell Differentiation Antigen.</i> Garland Publishing Inc.</li> <li>Diessner J, <i>et al.</i> 2014. <i>Cell Death Dis.</i> 5:1149. <u>PubMed</u></li> </ol>
Description:	CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from a low or mid level of intensity to high expression levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. CD44 is an adhesion molecule involved in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation.
Antigen References:	<ol> <li>Barclay AN, <i>et al.</i> 1997. The Leukocyte Antigen FactsBook Academic Press.</li> <li>Haynes BF, <i>et al.</i> 1991. <i>Cancer Cells</i> 3:347.</li> <li>Goldstein LA, <i>et al.</i> 1989. <i>Cell</i> 56:1063.</li> </ol>

4. Mikecz K, et al

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products. Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com