Alexa Fluor[®] 488 anti-human CD44

CD34 (dione f18) Aleva Plante 48		
	Human peripl lymphocytes	
	CD3 APC and (clone BJ18) A (left), or mou	
	Fluor® 488 is	

CDI VAC

Human peripheral blood lymphocytes were stained with CD3 APC and anti-human CD44 (clone BJ18) Alexa Fluor® 488 (left), or mouse IgG1, κ Alexa Fluor® 488 isotype control (right).

Applications:

Applications: Flow Cytometry Recommended Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the Usage: suggested use of this reagent is 5 μ l per million cells in 100 μ l staining volume or 5 μ l per 100 μ l of whole blood. * Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at 488 nm. Application 1. Kishimoto T, et al. eds. 1997 Leucocyte Typing VI:White Cell **References:** Differentiation Antigen. Garland Publishing Inc. **Description:** CD44 is a 80-95 kD glycoprotein also known as Hermes, Pgp1, H-CAM, or HUTCH. It is expressed on all leukocytes, endothelial cells, hepatocytes, and mesenchymal cells. As B and T cells become activated or progress to the memory stage, CD44 expression increases from a low or mid level of intensity to high expression levels. Thus, CD44 has been reported to be a valuable marker for memory cell subsets. CD44 is an adhesion molecule involved in leukocyte attachment to and rolling on endothelial cells, homing to peripheral lymphoid organs and to the sites of inflammation, and leukocyte aggregation. Antigen 1. Barclay AN, et al. 1997. The Leukocyte Antigen FactsBook Academic **References:** Press. 2. Haynes BF, et al. 1991. Cancer Cells 3:347. 3. Goldstein LA, et al. 1989. Cell 56:1063. 4. Mikecz K, et al. 1995. Nat. Med. 1:558.