

APC anti-human CD122 (IL-2R β)

Catalog # / Size: 2295040 / 100 tests
2295035 / 25 tests

Clone: TU27

Isotype: Mouse IgG1, κ

Immunogen: TL-Mor cell line


Reactivity: Human

Preparation: The antibody was conjugated with APC under optimal conditions, and is at >85% purity. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Workshop Number: V C050

Concentration: Lot-specific



Human peripheral blood lymphocytes stained with CD56 (HCD56) PE and TU27 APC

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications include (for the relevant formats) include: immunoprecipitation, blocking of IL-2 binding to CD122, and partial inhibition of IL-2 induced cell proliferation.

Application References: 1. Takeshita T, *et al.* 1989. *J. Exp. Med.* 169:1323.
2. Marra P, *et al.* 2014. *Cancer Res.* 74:4908. [PubMed](#)

Description: CD122 is a 70-75 kD type I transmembrane glycoprotein and member of the Ig superfamily. It is IL-2 receptor β chain also known as IL-2R β , which is also shared by the IL-15 receptor. CD122 is constitutively expressed by NK cells and at lower levels by a subset of T cells. Its expression is upregulated upon activation. The IL-2R β chain can combine with either the common γ subunit (γ c, CD132) alone or with the γ c subunit and the IL-2R α subunit (CD25) to generate intermediate or high affinity IL-2 receptor complexes, respectively. CD122 expression levels can be upregulated by activation.

Antigen References: 1. Zola H, *et al.* 2007. *Leukocyte and Stromal Cell Molecules: The CD Markers* Wiley-Liss A John Wiley & Sons Inc, Publication
2. Minami Y, *et al.* 1993. *Annu. Rev. Immunol.* 11:245.
3. Suzuki H, *et al.*