

**PerCP/Cy5.5 anti-human CD122 (IL-2R $\beta$ )**

**Catalog # / Size:** 2295055 / 25 tests  
2295060 / 100 tests

**Clone:** TU27

**Isotype:** Mouse IgG1,  $\kappa$

**Immunogen:** TL-Mor cell line


**Reactivity:** Human

**Preparation:** The antibody was purified by affinity chromatography and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated PerCP/Cy5.5 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Workshop Number:** V C050

**Concentration:** Lot-specific



Human peripheral blood lymphocytes were stained with CD56 FITC and CD122 (clone TU27) PerCP/Cy5.5 (top) or mouse IgG1,  $\kappa$  PerCP/Cy5.5 isotype control (bottom).

**Applications:**


**Applications:** Flow Cytometry

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

\* PerCP/Cy5.5 has a maximum absorption of 482 nm and a maximum emission of 690 nm.

**Application Notes:** Additional reported applications include (for the relevant formats) include: immunoprecipitation, blocking of IL-2 binding to CD122, and partial inhibition of IL-2 induced cell proliferation.

**Application References:** 1. Takeshita T, *et al.* 1989. *J. Exp. Med.* 169:1323.



**Description:** CD122 is a 70-75 kD type I transmembrane glycoprotein and member of the Ig superfamily. It is IL-2 receptor  $\beta$  chain also known as IL-2R $\beta$ , which is also shared by the IL-15 receptor. CD122 is constitutively expressed by NK cells and at lower levels by a subset of T cells. Its expression is upregulated upon activation. The IL-2R $\beta$  chain can combine with either the common  $\gamma$  subunit ( $\gamma$ c, CD132) alone or

with the  $\gamma_c$  subunit and the IL-2R $\alpha$  subunit (CD25) to generate intermediate or high affinity IL-2 receptor complexes, respectively. CD122 expression levels can be upregulated by activation.

**Antigen  
References:**

1. Zola H, *et al.* 2007. *Leukocyte and Stromal Cell Molecules*:The CD Markers Wiley-Liss A John Wiley & Sons Inc, Publication
2. Minami Y, *et al.* 1993. *Annu. Rev. Immunol.* 11:245.
3. Suzuki H, *et al*