PE/Cy7 anti-human CD122 (IL-2Rβ)

Catalog # / Size: 2295065 / 25 tests

2295070 / 100 tests

Clone: TU27

Isotype: Mouse IgG1, κ

Immunogen: TL-Mor cell line

Reactivity: Human

Preparation: The antibody was purified by affinity

chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7

and unconjugated antibody.


Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Workshop Number: V C050

Concentration: Lot-specific



Human peripheral blood lymphocytes were stained with CD56 APC and CD122 (clone TU27) PE/Cy7 (top) or mouse IgG1, κ PE/Cy7 isotype control (bottom).

CD56 APC

Applications:

Usage:

Applications: Flow Cytometry

Recommended Each lot of this antibody is quality

control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is

recommended that the reagent be titrated for optimal performance for

each application.

Application Additional reported applications include **Notes:** (for the relevant formats) include:

(for the relevant formats) include: immunoprecipitation, blocking of IL-2 binding to CD122, and partial inhibition

of IL-2 induced cell proliferation.

Application References:

1. Takeshita T, et al. 1989. J. Exp. Med. 169:1323.

Description: CD122 is a 70-75 kD type I transmembrane glycoprotein and member of the Ig

superfamily. It is IL-2 receptor β chain also known as IL-2R β , which is also shared by the IL-15 receptor. CD122 is constitutively expressed by NK cells and at lower levels by a subset of T cells. Its expression is upregulated upon activation. The IL-2R β chain can combine with either the common γ subunit (γ c, CD132) alone or with the γ c subunit and the IL-2R α subunit (CD25) to generate intermediate or high affinity IL-2 receptor complexes, respectively. CD122 expression levels can

Nouse IgG1, x PE/Cy7

be upregulated by activation.

Antigen 1. Zola H, et al. 2007. Leukocyte and Stromal Cell Molecules: The CD Markers

Wiley-Liss A John Wiley & Sons Inc, Publication 2. Minami Y, et al. 1993. Annu. Rev. Immunol. 11:245. **References:** 3. Suzuki H, et al