SONY

Product Data Sheet

FITC anti-human CD158 (KIR2DL1/S1/S3/S5)

Catalog # / Size: 2297515 / 25 tests
2297520 / 100 tests

Clone: HP-MA4

Isotype: Mouse IgG2b, κ

Immunogen: Human NK cell clone LB2

Reactivity: Human, Non-human primate

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and BSA (origin USA).

Workshop Number: V T-072

Concentration: Lot-specific



Human peripheral blood lymphocytes stained with HP-MA4 FITC and CD56 APC

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 µl to 5 µl per test**. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 µl staining volume or per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: mAb HP-MA4 reacts with KIR2DL1 (CD158a), KIR2DS1 (CD158h), KIR2DS3, and KIR2DS5 (CD158g).

Additional reported applications include: inhibits NK cell mediated cytotoxicity and immunoprecipitation.



Human peripheral blood lymphocytes were stained with anti-human CD50 (clone CBR-IC3/1) PerCP/Cy5.5 (closed histogram) or mouse IgG1

Application References:

1. De Miguel M and M. Lopez-Botet. 2002. *Inmunologia.* 21:187
2. Goodridge JP, *et al.* 2013. *J. Immunol.* 191:3553. PubMed

Description: CD158 molecules, also known as KIRs (killer cell immunoglobulin-like receptors), are a family of transmembrane proteins with either two (KIR2D) or three (KIR3D) Ig-like extracellular domains. Some KIRs with long cytoplasmic domains contain ITIMs and posses inhibitory functions and others with short cytoplasmic region lack ITIM and have activation functions. 14 polymorphic KIR genes have been reported in humans. CD158 is mainly expressed on a subset of NK cells and a small population of $CD8^+$ T cells. HLA-C is the ligand of CD158a/h.

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.

Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com

Antigen References: 1. Zola H, *et al.* eds. 2007. *Leukocyte and Stromal Cell Molecules:The CD Markers.* Wiely-Liss A John Wiley & Sons Inc, Publication

For research use only. Not for diagnostic use. Not for resale. Sony Biotechnology Inc. will not be held responsible for patent infringement or other violations that may occur with the use of our products.
Sony Biotechnology Inc. 1730 North First Street, San Jose, CA 95112 www.sonybiotechnology.com