## FITC anti-human CD158 (KIR2DL1/S1/S3/S5)

Catalog # / Size: 2297520 / 100 tests

2297515 / 25 tests

Clone: HP-MA4

**Isotype:** Mouse IgG2b, κ

Immunogen: Human NK cell clone LB2

Reactivity: Human

**Preparation:** The antibody was purified by affinity


chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2,

containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific



Human peripheral blood

lymphocytes stained with HP-MA4

FITC and CD56 APC

## **Applications:**

**Applications:** Flow Cytometry

Recommended

Usage:

Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 microL to 5 microL per test**. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 microL staining volume or per 100 microL of whole blood. It is recommended that the reagent be titrated for

optimal performance for each application.

**Application** 

Notes: KIR2DS5 (

mAb HP-MA4 reacts with KIR2DL1 (CD158a), KIR2DS1 (CD158h), KIR2DS3, and

KIR2DS5 (CD158g).

Additional reported applications include: inhibits NK cell mediated cytotoxicity

and immunoprecipitation.

Application References:

1. De Miguel M and M. Lopez-Botet. 2002. Inmunologia. 21:187

2. Goodridge JP, et al. 2013. J. Immunol. 191:3553. PubMed

**Description:** CD158 molecules, also known as KIRs (killer cell immunoglobulin-like receptors),

are a family of transmembrane proteins with either two (KIR2D) or three (KIR3D) Ig-like extracellular domains. Some KIRs with long cytoplasmic domains contain ITIMs and posses inhibitory functions and others with short cytoplasmic region lack ITIM and have activation functions. 14 polymorphic KIR genes have been reported in humans. CD158 is mainly expressed on a subset of NK cells and a

small population of CD8<sup>+</sup> T cells. HLA-C is the ligand of CD158a/h.

Antigen

1. Zola H, et al. eds. 2007. Leukocyte and Stromal Cell Molecules: The CD Markers.

**References:** Wiely-Liss A John Wiley & Sons Inc, Publication