

PerCP anti-human CD161

Catalog # / Size: 2299665 / 25 tests
2299670 / 100 tests

Clone: HP-3G10

Isotype: Mouse IgG1, κ


Immunogen: Human NK cells

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography and conjugated with PerCP under optimal conditions. The solution is free of unconjugated PerCP and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Concentration: Lot-specific




Human peripheral blood lymphocytes were stained with CD4 FITC and CD161 (clone HP-3G10) PerCP (top) or mouse IgG1, κ PerCP isotype control (bottom).

Applications:

Applications: Flow Cytometry

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For flow cytometric staining, the suggested use of this reagent is 5 microL per million cells or 5 microL per 100 microL of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.



* PerCP has a maximum absorption of 482 nm and a maximum emission of 675 nm.

Application Notes: Additional reported applications (for the relevant formats) include: inhibition of cytokine production and Western blotting under nonreducing conditions.

Application References:

1. Gumá M, et al. 2004. *Blood* 104:3664.
2. Exley M, et al. 1998. *J. Exp. Med.* 188:867.
3. Marquez C, et al. 1998. *Blood* 91:2760.

Description: CD161 is a type II transmembrane glycoprotein, also known as NKR-P1A, that is expressed as a 40-44 kD homodimer. It is a member of the C-type lectin superfamily. CD161 is expressed on a majority of NK cells, NKT cells, and subsets of peripheral T cells and CD3⁺ thymocytes. It has been reported that Th17 cells are a subpopulation of CD4⁺CD161⁺CCR6⁺ cells. While the biological function of CD161 is not clear, it has been suggested to serve either as a stimulatory

receptor or to inhibit NK cell-mediated cytotoxicity and cytokine production. LLT-1 (lectin-like transcript-1, also named as osteoclast inhibitory lectin or CLEC2D) is the ligand of CD161.

**Antigen
References:**

1. Takahashi T, *et al.* 2006. *J. Immunol.* 176:211.
2. Cosmi L, *et al.* 2008. *J. Exp. Med.* 205:1903.
3. Aldemir H, *et al.* 2005. *J. Immunol.* 175:7791.
4. Rosen DB, *et al.* 2008. <